arXiv:1105.4398v3 [math.RA] 5 Apr 2012

SEMISIMPLE HOPF ALGEBRAS OF DIMENSION $2q^3$

JINGCHENG DONG AND LI DAI

Abstract. Let q be a prime number, k an algebraically closed field of characteristic 0, and H a non-trivial semisimple Hopf algebra of dimension $2q^3$. This paper proves that H can be constructed either from group algebras and their duals by means of extensions, or from Radford's biproduct $H \cong R\#kG$, where kG is the group algebra of G of order 2, R is a semisimple Yetter-Drinfeld Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$ of dimension q^3.

1. Introduction

In the last twenty years, various classification results were obtained for finite-dimensional semisimple Hopf algebras over an algebraically closed field of characteristic 0. Up to now, semisimple Hopf algebras of dimension p, p^2, p^3, pq, pq^2 and pqr have been completely classified, where p, q, r are distinct prime numbers. See [3, 4, 8, 9, 22] for details.

In the present paper, we shall continue the investigation on the classification of semisimple Hopf algebras. The main purpose of this paper is to investigate semisimple Hopf algebras of dimension $2q^3$, where q is a prime number. Of course, some other interesting results are also obtained in this paper.

The paper is organized as follows. In Section 2, we recall the definitions and basic properties of semisolvability, characters, Radford's biproducts and Drinfeld double, respectively. Some useful lemmas are also contained in this section. In Section 3, we study the structure of non-trivial semisimple Hopf algebras of dimension pq^3, where p, q are prime numbers with $p^2 < q$. In Section 4, we study the structure of non-trivial semisimple Hopf algebras of dimension $2q^3$, where q is a prime number.

Throughout this paper, all modules and comodules are left modules and left comodules, and moreover they are finite-dimensional over an algebraically closed field k of characteristic 0. $\otimes$, dim mean $\otimes_k$, $\dim_k$, respectively. For two positive integers m and n, $gcd(m, n)$ denotes the greatest common divisor of m, n. Our references for the theory of Hopf algebras are [11] or [21]. The notation for Hopf algebras is standard. For example, the group of group-like elements in H is denoted by $G(H)$.

2. Preliminaries

2.1. **Characters.** Throughout this subsection, H will be a semisimple Hopf algebra over k.

Let V be an H-module. The character of V is the element $\chi = \chi_V \in H^*$ defined by $\langle \chi, h\rangle = \mathrm{Tr}_V(h)$ for all $h \in H$. The degree of χ is defined to be the

---

2000 *Mathematics Subject Classification.* 16W30.

Key words and phrases. semisimple Hopf algebra, semisolvability, Radford's biproduct, character, Drinfeld double.

integer $\deg\chi = \chi(1) = \dim V$. We shall use X_t to denote the set of all irreducible characters of H of degree t. All irreducible characters of H span a subalgebra $R(H)$ of H^*, which is called the character algebra of H. The antipode S induces an anti-algebra involution $* : R(H) \to R(H)$, given by $\chi \mapsto \chi^* := S(\chi)$. The character of the trivial H-module is the counit ε.

Let $\chi_U, \chi_V \in R(H)$ be the characters of the H-modules U and V, respectively. The integer $m(\chi_U, \chi_V) = \dim\mathrm{Hom}_H(U, V)$ is defined to be the multiplicity of U in V. Let $\widehat{H}$ denote the set of irreducible characters of H. Then $\widehat{H}$ is a basis of $R(H)$. If $\chi \in R(H)$, we may write $\chi = \sum_{\alpha\in\widehat{H}} m(\alpha, \chi)\alpha$.

For any group-like element g in $G(H^*)$, $m(g, \chi\chi^*) > 0$ if and only if $m(g, \chi\chi^*) = 1$ if and only if $g\chi = \chi$. The set of such group-like elements forms a subgroup of $G(H^*)$, of order at most $(\deg\chi)^2$. See [15, Theorem 10]. Denote this subgroup by $G[\chi]$. In particular, we have

$$\chi\chi^* = \sum_{g\in G[\chi]} g + \sum_{\alpha\in\widehat{H}, \deg\alpha>1} m(\alpha, \chi\chi^*)\alpha. \tag{2.1}$$

A subalgebra A of $R(H)$ is called a standard subalgebra if A is spanned by irreducible characters of H. Let X be a subset of $\widehat{H}$. Then X spans a standard subalgebra of $R(H)$ if and only if the product of characters in X decomposes as a sum of characters in X. There is a bijection between $*$-invariant standard subalgebras of $R(H)$ and quotient Hopf algebras of H. See [15, Theorem 6].

H is said to be of type $(d_1, n_1; \cdots ; d_s, n_s)$ as an algebra if $d_1 = 1, d_2, \cdots , d_s$ are the dimensions of the simple H-modules and n_i is the number of the non-isomorphic simple H-modules of dimension d_i. That is, as an algebra, H is isomorphic to a direct product of full matrix algebras

$$H \cong k^{(n_1)} \times \prod_{i=2}^{s} M_{d_i}(k)^{(n_i)}.$$

If H^* is of type $(d_1, n_1; \cdots ; d_s, n_s)$ as an algebra, then H is said to be of type $(d_1, n_1; \cdots ; d_s, n_s)$ as a coalgebra.

Lemma 2.1. *Let χ be an irreducible character of H. Then*

(1) The order of $G[\chi]$ divides $(\deg\chi)^2$.

(2) The order of $G(H^*)$ divides $n(\deg\chi)^2$, where n is the number of non-isomorphic irreducible characters of degree $\deg\chi$.

Proof. It follows from Nichols-Zoeller Theorem [16]. See also [14, Lemma 2.2.2]. □

2.2. **Semisolvability.** Let B be a finite-dimensional Hopf algebra over k. A Hopf subalgebra $A \subseteq B$ is called normal if $h_1AS(h_2) \subseteq A$ and $S(h_1)Ah_2 \subseteq A$, for all $h \in B$. If B does not contain proper normal Hopf subalgebras then it is called simple. The notion of simplicity is self-dual, that is, B is simple if and only if B^* is simple.

Let $\pi : H \to B$ be a Hopf algebra map and consider the subspaces of coinvariants

$$H^{co\pi} = \{h \in H | (id \otimes \pi)\Delta(h) = h \otimes 1\}, \text{and}$$
$${}^{co\pi}H = \{h \in H | (\pi \otimes id)\Delta(h) = 1 \otimes h\}.$$

Then $H^{co\pi}$ (respectively, ${}^{co\pi}H$) is a left (respectively, right) coideal subalgebra of H. Moreover, we have

$$\dim H = \dim H^{co\pi} \dim\pi(H) = \dim {}^{co\pi}H \dim\pi(H).$$

The left coideal subalgebra $H^{co\pi}$ is stable under the left adjoint action of H. Moreover $H^{co\pi} = {}^{co\pi}H$ if and only if $H^{co\pi}$ is a (normal) Hopf subalgebra of H. If this is the case, we shall say that the map $\pi : H \to B$ is normal. See [19] for more details.

The following lemma comes from [13, Section 1.3].

Lemma 2.2. *Let $\pi : H \to B$ be a Hopf epimorphism and A a Hopf subalgebra of H such that $A \subseteq H^{co\pi}$. Then* $\dim A$ *divides* $\dim H^{co\pi}$.

The notions of upper and lower semisolvability for finite-dimensional Hopf algebras have been introduced in [10], as generalizations of the notion of solvability for finite groups. By definition, H is called lower semisolvable if there exists a chain of Hopf subalgebras

$$H_{n+1} = k \subseteq H_n \subseteq \cdots \subseteq H_1 = H$$

such that H_{i+1} is a normal Hopf subalgebra of H_i, for all i, and all quotients $H_i/H_iH_{i+1}^+$ are trivial. That is, they are isomorphic to a group algebra or a dual group algebra. Dually, H is called upper semisolvable if there exists a chain of quotient Hopf algebras

$$H_{(0)} = H \xrightarrow{\pi_1} H_{(1)} \xrightarrow{\pi_2} \cdots \xrightarrow{\pi_n} H(n) = k$$

such that $H_{(i-1)}^{co\pi_i}$ is a normal Hopf subalgebra of $H_{(i-1)}$, and all $H_{(i-1)}^{co\pi_i}$ are trivial.

In analogy with the situations for finite groups, it is enough for many applications to know that a Hopf algebra is semisolvable.

By [10, Corollary 3.3], we have that H is upper semisolvable if and only if H^* is lower semisolvable. If this is the case, then H can be obtained from group algebras and their duals by means of (a finite number of) extensions.

Proposition 2.3. *Let H be a semisimple Hopf algebra of dimension pq^3, where p, q are distinct prime numbers. If H is not simple as a Hopf algebra then it is semisolvable.*

Proof. By assumption, H has a proper normal Hopf subalgebra K. Moreover, by Nichols-Zoeller Theorem [16], $\dim K$ divides $\dim H = pq^3$. We shall examine every possible $\dim K$.

If $\dim K = q^2$ or pq then $k \subseteq K \subseteq H$ is a chain such that K and H/HK^+ are both trivial (see [3, 8]). Hence, H is lower semisolvable.

If $\dim K = q^3$ then [8] shows that K has a non-trivial central group-like element g. Let $L = k\langle g\rangle$ be the group algebra of the cyclic group $\langle g\rangle$ generated by g. Then $k \subseteq L \subseteq K \subseteq H$ is a chain such that $L, K/KL^+$ and H/HK^+ are all trivial (see [22]). Hence, H is lower semisolvable.

If $\dim K = pq^2$ then [2, Lemma 2.2] and [12, Theorem 5.4.1] show that K has a proper normal Hopf subalgebra L of dimension p, q, pq or q^2. Then $k \subseteq L \subseteq K \subseteq H$ is a chain such that $L, K/KL^+$ and H/HK^+ are all trivial. Hence, H is lower semisolvable.

Finally, we consider the case that $\dim K = p$ or q. Let L be a proper normal Hopf subalgebra of H/HK^+ (Notice that H/HK^+ is not simple). Write $\overline{K} = H/HK^+$ and $\overline{L} = \overline{K}/\overline{K}L^+$. Then $H \xrightarrow{\pi_1} \overline{K} \xrightarrow{\pi_2} \overline{L} \to k$ is a chain such that every map is normal and $H^{co\pi_1}$, $(\overline{K})^{co\pi_2}$ are trivial. Hence, H is upper semisolvable. □

2.3. **Radford's biproduct.** Let A be a semisimple Hopf algebra and let ${}^A_A\mathcal{YD}$ denote the braided category of Yetter-Drinfeld modules over A. Let R be a semisimple Yetter-Drinfeld Hopf algebra in ${}^A_A\mathcal{YD}$. Denote by $\rho : R \to A \otimes R$, $\rho(a) = a_{-1} \otimes a_0$, and $\cdot : A \otimes R \to R$, the coaction and action of A on R, respectively. We shall use the notation $\Delta(a) = a^1 \otimes a^2$ and S_R for the comultiplication and the antipode of R, respectively.

Since R is in particular a module algebra over A, we can form the smash product (see [11, Definition 4.1.3]). This is an algebra with underlying vector space $R \otimes A$, multiplication is given by

$$(a \otimes g)(b \otimes h) = a(g_1 \cdot b) \otimes g_2 h, \text{ for all } g, h \in A, a, b \in R,$$

and unit $1 = 1_R \otimes 1_A$.

Since R is also a comodule coalgebra over A, we can dually form the smash coproduct. This is a coalgebra with underlying vector space $R \otimes A$, comultiplication is given by

$$\Delta(a \otimes g) = a^1 \otimes (a^2)_{-1} g_1 \otimes (a^2)_0 \otimes g_2, \text{ for all } h \in A, a \in R,$$

and counit $\varepsilon_R \otimes \varepsilon_A$.

As observed by D. E. Radford (see [17, Theorem 1]), the Yetter-Drinfeld condition assures that $R \otimes A$ becomes a Hopf algebra with these structures. This Hopf algebra is called the Radford's biproduct of R and A. We denote this Hopf algebra by $R\#A$ and write $a\#g = a \otimes g$ for all $g \in A, a \in R$. Its antipode is given by

$$S(a\#g) = (1\#S(a_{-1}g))(S_R(a_0)\#1), \text{ for all } g \in A, a \in R.$$

A biproduct $R\#A$ as described above is characterized by the following property(see [17, Theorem 3]): suppose that H is a finite-dimensional Hopf algebra endowed with Hopf algebra maps $\iota : A \to H$ and $\pi : H \to A$ such that $\pi\iota : A \to A$ is an isomorphism. Then the subalgebra $R = H^{co\pi}$ has a natural structure of Yetter-Drinfeld Hopf algebra over A such that the multiplication map $R\#A \to H$ induces an isomorphism of Hopf algebras.

Following [20, Proposition 1.6], $H \cong R\#A$ is a biproduct if and only if $H^* \cong R^*\#A^*$ is a biproduct.

The following lemma is a special case of [13, Lemma 4.1.9].

Lemma 2.4. *Let H be a semisimple Hopf algebra of dimension pq^3, where p, q are distinct prime numbers. If $gcd(|G(H)|, |G(H^*)|) = p$, then $H \cong R\#kG$ is a biproduct, where kG is the group algebra of group G of order p, R is a semisimple Yetter-Drinfeld Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$ of dimension q^3.*

2.4. **Drinfeld double.** For a finite-dimensional Hopf algebra H, $D(H) = H^{*cop} \bowtie H$ will denote the Drinfeld double of H. $D(H)$ is a Hopf algebra with underlying vector space $H^{*cop} \otimes H$. More details on $D(H)$ can be found in [11, Section 10.3]. The following theorem follows directly from [18, Proposition 9,10].

Theorem 2.5. *Suppose that H is a finite-dimensional Hopf algebra.*

(1) The map $G(H^*) \times G(H) \to G(D(H))$, given by $(\eta, g) \mapsto \eta \bowtie g$, is a group isomorphism.

(2) Every group-like element of $D(H)^*$ is of the form $g \otimes \eta$, where $g \in G(H)$ and $\eta \in G(H^*)$. Moreover, $g \otimes \eta \in G(D(H)^*)$ if and only if $\eta \bowtie g$ is in the center of $D(H)$.

Corollary 2.6. *Suppose that* H *is a finite-dimensional Hopf algebra such that* $G(D(H)^*)$ *is non-trivial. If* $gcd(|G(H)|, |G(H^*)|) = 1$ *then* H *or* H^* *has a non-trivial central group-like element.*

Proof. Let $1 \neq g \otimes \eta \in G(D(H)^*)$. We may assume that $1 \neq g \in G(H)$, since otherwise $\eta \in G(H^*)$ would be a non-trivial central group-like element, and similarly we may assume that $\varepsilon \neq \eta \in G(H^*)$. Since $gcd(|G(H)|, |G(H^*)|) = 1$, the order of g and η are different. Assume that the order of g is n. Then $(g \otimes \eta)^n = g^n \otimes \eta^n = 1 \otimes \eta^n \neq 1 \otimes \varepsilon$ implies that $\eta^n \bowtie 1$ is in the center of $D(H)$. Hence, η^n is a non-trivial central group-like element in $G(H^*)$. Similarly, we can prove that $G(H)$ also has a non-trivial central group-like element. $\square$

3. Semisimple Hopf algebras of dimension pq^3

Lemma 3.1. *Let* H *be a semisimple Hopf algebra of dimension* pq^3*, where* $p < q$ *are prime numbers. If* H *has a Hopf subalgebra* K *of dimension* q^3 *then* H *is lower semisolvable.*

Proof. Since the index $[H : K] = p$ is the smallest prime number dividing $\dim H$, the result in [7] shows that K is a normal Hopf algebra of H. The lemma then follows from Proposition 2.3. $\square$

In the rest of this section, p, q will be distinct prime numbers such that $p^2 < q$, and H will be a non-trivial semisimple Hopf algebra of dimension pq^3.

Recall that a semisimple Hopf algebra A is called of Frobenius type if the dimensions of the simple A-modules divide the dimension of A. Kaplansky conjectured that every finite-dimensional semisimple Hopf algebra is of Frobenius type [5, Appendix 2]. It is still an open problem. However, many examples show that a positive answer to Kaplansky's conjecture would be very helpful in the classification of semisimple Hopf algebras.

By [2, Lemma 2.2], H is of Frobenius type and $|G(H^*)| \neq 1$. Therefore, the dimension of a simple H-module can only be $1, p, q$ or pq. It follows that we have an equation

$$pq^3 = |G(H^*)| + p^2a + q^2b + p^2q^2c, \tag{3.1}$$

where a, b, c are the numbers of non-isomorphic simple H-modules of dimension p, q and pq, respectively. By Nichols-Zoeller Theorem [16], the order of $G(H^*)$ divides $\dim H$. We shall give some results concerning the order of $G(H^*)$.

Lemma 3.2. *The order of* $G(H^*)$ *can not be* q.

Proof. Suppose on the contrary that $|G(H^*)| = q$. Let χ be an irreducible character of degree p. By Lemma 2.1 (1) and the fact that $G[\chi]$ is a subgroup of $G(H^*)$, we know that $G[\chi] = \{\varepsilon\}$ is trivial. It follows that the decomposition of $\chi\chi^*$ (2.1) gives rise to a contradiction, since $p^2 < q$. Therefore, $a = 0$ and equation (3.1) is $pq^3 = q + q^2b + p^2q^2c$, which is impossible. $\square$

Lemma 3.3. *If* $|G(H^*)| = q^2$ *then* $a = 0$ *and* $b \neq 0$.

Proof. A similar argument as in Lemma 3.2 shows that $a = 0$. Therefore, equation (3.1) is $pq^3 = q^2 + q^2b + p^2q^2c$. Obviously, $b \neq 0$, otherwise a contradiction will occur. $\square$

Lemma 3.4. *If* $|G(H^*)| = q^3$ *then* H *is upper semisolvable.*

Proof. It follows from Lemma 3.1. □

Lemma 3.5. *If* $|G(H^*)| = q^2$ *and* H *has a Hopf subalgebra* K *of dimension* pq^2 *then* H *is lower semisolvable.*

Proof. Considering the map $\pi : H^* \to K^*$ obtained by transposing the inclusion $K \subseteq H$, we have $\dim(H^*)^{co\pi} = q$ (see Section 2.2). By Lemma 3.3, the dimension of every irreducible left coideal of H^* is $1, q$ or pq. Therefore, by Lemma 2.2, as a left coideal of H^*, $(H^*)^{co\pi}$ decomposes in the form $(H^*)^{co\pi} = k\langle g\rangle$, where $\langle g\rangle$ is a subgroup of $G(H^*)$ generated by g which is of order q. Therefore, $(H^*)^{co\pi}$ is a normal Hopf subalgebra of H^*. The lemma then follows from Proposition 2.3. □

Lemma 3.6. *If* $|G(H^*)| = p$ *then* H *is either semisolvable, or isomorphic to a Radford's biproduct* $H \cong R\#kG$*, where* kG *is the group algebra of* G *of order* p*,* R *is a semisimple Yetter-Drinfeld Hopf algebra in* ${}^{kG}_{kG}\mathcal{YD}$ *of dimension* q^3*.*

Proof. Notice that $G(D(H)^*)$ is not trivial by [2, Lemma 2.2]. If $|G(H)| = q^2$ or q^3 then H is not simple by Corollary 2.6. Hence, H is semisolvable. If $|G(H)| = p, pq$ or pq^2 then, by Lemma 2.4, H is isomorphic to a Radford's biproduct $H \cong R\#kG$, where kG is the group algebra of G of order p, R is a semisimple Yetter-Drinfeld Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$ of dimension q^3. □

4. SEMISIMPLE HOPF ALGEBRAS OF DIMENSION $2q^3$

In this section, H will be a non-trivial semisimple Hopf algebra of dimension $2q^3$, where q is a prime number. We shall discuss the structure of H. When $q = 2$, the structure of H is given in [6]. When $q = 3$, the structure of H is given in [13, Chapter 12]. Therefore, in the rest of this section, we always assume that $q > 3$.

Using notations from Section 3, we have

$$2q^3 = |G(H^*)| + 4a + q^2b + 4q^2c. \tag{4.1}$$

By the results in Section 3, it suffices to consider the cases that $|G(H^*)| = 2q, 2q^2$ and q^2.

Lemma 4.1. *If* $|G(H^*)| = 2q$ *or* $2q^2$ *then* H *is either semisolvable, or isomorphic to a Radford's biproduct* $H \cong R\#kG$*, where* kG *is the group algebra of* G *of order* 2*,* R *is a semisimple Yetter-Drinfeld Hopf algebra in* ${}^{kG}_{kG}\mathcal{YD}$ *of dimension* q^3*.*

Proof. By Lemma 3.4 and 3.6, we may assume that $|G(H)| \neq 2$ and q^3.

First, if $gcd(|G(H)|, |G(H^*)|) = 2$ then the lemma follows from Lemma 2.4.

Second, we assume that $|G(H^*)| = 2q$ and $|G(H)| = q^2$. Notice that $a \neq 0$, otherwise equation (4.1) will give rise to a contradiction. Hence, $G(H^*) \cup X_2$ spans a standard subalgebra of $R(H)$. It follows that H has a quotient Hopf algebra K of dimension $2q + 4a$. By Nichols-Zoeller Theorem [16], $\dim K = 2q^2$ or $2q^3$. If $\dim K = 2q^2$ then the duality of Lemma 3.5 proves the lemma. If $\dim K = 2q^3$ then H is of type $(1, 2q; 2, a)$ as an algebra. Then [1, Theorem 6.4] shows that either H or H^* must contain a non-trivial central group-like element. The lemma then follows from Proposition 2.3.

Finally, we assume that $|G(H^*)| = 2q^2$ and $|G(H)| = q^2$. In this case, the duality of Lemma 3.5 proves the lemma. □

Lemma 4.2. *If* $|G(H^*)| = q^2$ *then* H *is semisolvable.*

Proof. By the discussion above, we may assume that $|G(H)| = q^2$. Considering the map $\pi : H \to (kG(H^*))^*$ obtained by transposing the inclusion $kG(H^*) \subseteq H^*$, we have $\dim H^{co\pi} = 2q$. By Lemma 3.3, the dimension of every irreducible left coideal of H is $1, q$ or $2q$. Therefore, by Lemma 2.2, as a left coideal of H, $H^{co\pi}$ decomposes in the form $H^{co\pi} = k\langle g\rangle \oplus V$, where $k\langle g\rangle$ is the group algebra of a subgroup $\langle g\rangle$ of $G(H)$ generated by g which is of order q, and V is an irreducible left coideal of H of dimension q. Since gV and Vg are irreducible left coideals of H isomorphic to V , and gV, Vg are contained in $H^{co\pi}$, we have $gV = V = Vg$. Then [13, Corollary 3.5.2] shows that $k\langle g\rangle$ is a normal Hopf subalgebra of $k[C]$, where C is the simple subcoalgebra of H containing V, and $k[C]$ is a Hopf subalgebra of H generated by C as an algebra. Clearly, $\dim k[C] \geq q + q^2$. Moreover, by Nichols-Zoeller Theorem [16], $\dim k[C] = 2q^3, q^3$ or $2q^2$. If $\dim k[C] = 2q^3$ then $k[C] = H$ and $k\langle g\rangle$ is a normal Hopf subalgebra of H. The lemma then follows from Proposition 2.3. If $\dim k[C] = q^3$ then the lemma follows from Lemma 3.1. If $\dim k[C] = 2q^2$ then the lemma follows from Lemma 3.5. □

From the discussion in Section 3 and 4, we obtain the main theorem.

Theorem 4.3. *Let H will be a non-trivial semisimple Hopf algebra of dimension $2q^3$, where $q > 3$ is a prime number. Then*

(1) If $\gcd(|G(H)|, |G(H^*)|) = 2$ then H is isomorphic to a Radford's biproduct $H \cong R\#kG$, where kG is the group algebra of G of order 2, R is a semisimple Yetter-Drinfeld Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$ of dimension q^3.

(2) In all other cases, H is semisolvable.

As an immediate consequence of Theorem 4.3, we have the following result.

Corollary 4.4. *If H is simple as a Hopf algebra then H is isomorphic to a Radford's biproduct $H \cong R\#kG$, where kG is the group algebra of G of order 2, R is a semisimple Yetter-Drinfeld Hopf algebra in ${}^{kG}_{kG}\mathcal{YD}$ of dimension q^3.*

References

[1] J. Bichon, S. Natale, Hopf algebra deformations of binary polyhedral groups. Trans. Groups, DOI: 10.1007/s00031-011-9133-x.

[2] J. Dong, Structure of semisimple Hopf algebras of dimension p^2q^2. arXiv:1009.3541v2, to appear in Communications in Algebra.

[3] P. Etingof and S. Gelaki, Semisimple Hopf algebras of dimension pq are trivial. J. Algebra 210(2), 664–669 (1998).

[4] P. Etingof, D. Nikshych and V. Ostrik, Weakly group-theoretical and solvable fusion categories. Adv. Math. 226 (1), 176–505 (2011).

[5] I. Kaplansky, Bialgebras. Chicago, University of Chicago Press 1975.

[6] Y. Kashina , Classification of semisimple Hopf algebras of dimension 16, J. Algebra 232, 617–663(2000).

[7] T. Kobayashi and A. Masuoka, A result extended from groups to Hopf algebras. Tsukuba J. Math. 21(1), 55–58 (1997).

[8] A. Masuoka, The p^n theorem for semisimple Hopf algebras. Proc. Amer. Math. Soc. 124, 735–737 (1996).

[9] A. Masuoka, Self-dual Hopf algebras of dimension p^3 obtained by extension. J. Algebra 178, 791-806 (1995)

[10] S. Montgomery and S. Whiterspoon, Irreducible representations of crossed products. J. Pure Appl. Algebra 129, 315–326 (1998).

[11] S. Montgomery, Hopf algebras and their actions on rings. CBMS Reg. Conf. Ser. Math. 82. Providence. Amer. Math. Soc. 1993.

[12] S. Natale, On semisimple Hopf algebras of dimension pq^r. Algebras Represent. Theory 7 (2), 173–188 (2004).
[13] S. Natale, Semisolvability of semisimple Hopf algebras of low dimension. Mem. Amer. Math. Soc. 186 (2007).
[14] S. Natale, On semisimple Hopf algebras of dimension pq^2. J. Algebra 221(2), 242–278 (1999).
[15] W. D. Nichols and M. B. Richmond, The Grothendieck group of a Hopf algebra. J. Pure Appl. Algebra 106, 297–306(1996).
[16] W. D. Nichols and M. B. Zoelle, A Hopf algebra freeness theorem. Amer. J. Math. 111(2), 381–385 (1989).
[17] D. Radford, The structure of Hopf algebras with a projection. J. Algebra 92, 322–347 (1985).
[18] D. Radford, Minimal Quasitriangular Hopf algebras. J. Algebra, 157, 285–315 (1993).
[19] H.-J. Schneider, Normal basis and transitivity of crossed products for Hopf algebras. J. Algebra 152, 289–312 (1992).
[20] Y. Sommerhäuser, Yetter-Drinfeld Hopf algebras over groups of prime order, Lectures Notes in Math. 1789, Springer-Verlag (2002).
[21] M. E. Sweedler, Hopf Algebras. New York, Benjamin 1969.
[22] Y. Zhu, Hopf algebras of prime dimension. Internat. Math. Res. Notices 1, 53–59 (1994).

College of Engineering, Nanjing Agricultural University, Nanjing 210031, Jiangsu, People's Republic of China
E-mail address, J. Dong: `dongjc@njau.edu.cn`
E-mail address, L. Dai: `daili1980@njau.edu.cn`